arXiv:1211.2236v1 [cond-mat.str-el] 9 Nov 2012

Nonequilibrium thermal transport and its relation to linear response

C. Karrasch[1,2], R. Ilan[1], and J. E. Moore[1,2]
[1] *Department of Physics, University of California, Berkeley, California 95720, USA and*
[2] *Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

We study the real-time dynamics of spin chains driven out of thermal equilibrium by an initial temperature gradient $T_L \neq T_R$. We demonstrate that the nonequilibrium energy current saturates fast to a finite value if the linear-response thermal conductivity is infinite, i.e. if the Drude weight D is nonzero. Our data suggests that a nonintegrable dimerized chain might support such dissipationless transport ($D > 0$). We show that the steady-state value of the current for arbitrary $T_L \neq T_R$ is completely determined by the linear conductance. Inhomogeneous systems exhibiting different bulk parameters as well as Luttinger liquid boundary physics induced by single impurities are discussed shortly.

PACS numbers: 75.10.Pq,71.27.+a,05.60.Gg

Introduction — One-dimensional (1d) electronic systems are realized in carbon nanotubes and individual polymer molecules and provide an approximate description of strongly anisotropic 3d materials. It has been known for many years that 1d systems can support unusual correlated electron phenomena such as Luttinger liquid physics. However, electrical and thermal transport in real materials are usually not governed by the free low-energy Luttinger liquid fixed point but by an interplay between dangerously irrelevant operators scattering the currents and conserved quantities protecting them [1–3].

In order to connect to actual experiments, it is thus essential to study generic microscopic models. Over the last decades a significant number of works [2–11] investigated equilibrium charge (or spin) transport properties. In particular, the question whether or not so-called integrable models, which possess a complete set of local conserved quantities, can support dissipationless currents at finite temperature was addressed extensively. Less is known about the quantitative effects of integrability-breaking perturbations which are naturally present in any experimental system, and even the qualitative question whether the linear-response conductivity of a nonintegrable model can still be infinite is not answered conclusively [12]. While experimental measurements of thermal transport driven by a temperature gradient in quasi-1d spin systems already exist [13, 14], only a few works investigate this theoretically [8, 9, 15–20]. Studying nonequilibrium thermal (or charge) transport is complicated in general – one reason being that is not even clear whether the long-time dynamics can be described by a low-energy theory – and constitutes one of the most active areas of research in strongly correlated condensed matter physics [21–29, 32].

The primary goal of our work is to obtain quantitative results on steady-state energy flow both near and far from equilibrium and to understand the effects of integrability and correlations. This is motivated by the experiments listed above and by recent technical advances in dynamical simulations [11]. As a prototypical model we consider a XXZ spin-1/2 chain in presence of two perturbations (dimerization and a staggered magnetic field) which break integrability [30, 31]. We demonstrate that the nonequilibrium energy current driven by a temperature gradient $T_L \neq T_R$ relaxes fast to a finite steady-state value if the linear-response thermal conductivity is infinite, i.e. if the Drude weight D is nonzero. Our data indicates that the dimerized chain might support such dissipationless transport ($D > 0$) despite the fact that it is nonintegrable (D can be extracted from the asymptote of the equilibrium energy current correlation function [8], and we cannot exclude that the latter decays on a hidden large time scale). We show that for a large class of problems the steady-state current takes the functional form $f(T_L) - f(T_R)$. This implies that nonequilibrium thermal transport is entirely determined by linear response – f can simply be obtained by integration of the equilibrium conductance $\partial_T f$. We demonstrate that at low temperatures the gapless integrable XXZ chain as well as the quantum Ising model exhibit universal nonequilibrium behavior conjectured by conformal field theory [32, 33]. Finally, we study inhomogeneous systems featuring different bulk interactions as well as Luttinger liquid physics [34] induced by an impurity at the interface.

Thermal non-equilibrium setup — We aim at investigating the real-time dynamics of the energy current $\langle J_E(n,t)\rangle$ through a one-dimensional infinite lattice system driven out of equilibrium by an initial sharp temperature gradient $T_L \neq T_R$. Our main focus is to study the long-time behavior of $\langle J_E(n,t)\rangle$ and specifically the question how it relates to linear-response thermal transport properties. As a prototypical model, we consider a chain of interacting spin-1/2 degrees of freedom $S_n^{x,y,z}$ governed by local Hamiltonians

$$h_n = J_n\big(S_n^x S_{n+1}^x + S_n^y S_{n+1}^y + \Delta_n S_n^z S_{n+1}^z\big) + b_n(S_n^z - S_{n+1}^z)\,, \tag{1}$$

or equivalently spinless Fermions through a Jordan-Wigner transformation. By choosing the couplings J_n,





FIG. 1. Energy current flowing between two semi-infinite spin-1/2 chains which are initially in thermal equilibrium at different temperatures $T_{L,R}$ and coupled at time $t = 0$ and position $n = 0$. (a) Integrable XXZ chain with z-anisotropy Δ. The behavior in the gapped phase $\Delta > 1$ is similar. (b) Nonintegrable dimerized XXZ chain where the coupling on every second bond is reduced by λ. The system is gapped. (c) XXZ chain in presence of a staggered field b rendering the model nonintegrable. A gap opens for $b \approx 0.3$. Despite the fact that the local energy density $h(n, t)$ does not relax (Inset), the current saturates fast to a unique finite value except for $b > 0$. We attribute this to a finite linear-response thermal Drude weight of both the pure XXZ chain and the nonintegrable dimerized chain (see Figure 2). This indicates an intricate relation between nonequilibrium and linear thermal transport properties.

FIG. 2. Linear-response energy current correlation function whose long-time asymptote determines the Drude weight through Equation (5). For the integrable XXZ chain ($\lambda = 1$, $b = 0$), the global energy current $J_E = \sum_n J_E(n)$ is conserved; thus, $\langle J_E(t)J_E\rangle = \langle J_E(0)J_E\rangle$, and DMRG can be compared with exact Bethe ansatz results (symbols). (a,b) Nonintegrable dimerized chain. The current correlation function seems to saturate at a finite value (or decays on a hidden large time scale), indicating a finite Drude weight $D > 0$. (c) The data in presence of a staggered field $b > 0$ is consistent with $D = 0$.

Δ_n, and b_n appropriately:

$$J_n = \begin{cases} 1 & n \text{ odd} \\ \lambda & n \text{ even} \end{cases} , \quad \Delta_n = \Delta , \quad b_n = \frac{(-1)^n b}{2} , \qquad (2)$$

we can study systems which are gapless or gapped and – as a key aspect of this work – investigate the role of integrability. For $\lambda = 1$ and $b = 0$, Equation (1) can be diagonalized via Bethe ansatz [35]; the model is nonintegrable otherwise. The spectrum is gapless for $|\Delta| \leq 1$ and gapped for $\Delta > 1$. A gap opens for $\lambda < 1$ or sufficiently large $b > b_c$; $b_c > 0$ if $-1 < \Delta < -1/\sqrt{2}$ [31, 36].

Thermal nonequilibrium is introduced via the following protocol: We initially consider two seperate semi-infinite chains ($N \to \infty$), $H_0 = H_L + H_R = \sum_{n=-N/2+1}^{-1} h_n + \sum_1^{N/2-1} h_n$, each being in thermal (grand-canonical) equilibrium at temperatures T_L and T_R. The corresponding density matrix factorizes, $\rho_0 = \rho_L \otimes \rho_R$, $\rho_i = \exp(-H_i/T_i)/\mathrm{Tr}\exp(-H_i/T_i)$. At time $t = 0$, the chains are coupled through h_0, and the time evolution of ρ_0 is computed w.r.t. $H = H_0 + h_0$. The energy current is defined by a continuity equation [8],

$$\partial_t h_n = J_E(n) - J_E(n+1) \ \Rightarrow \ J_E(n) = i[h_{n-1}, h_n] , \quad (3)$$

and its time evolution is simply given by

$$\langle J_E(n,t)\rangle = \mathrm{Tr}\left[e^{iHt}\rho_0 e^{-iHt} J_E(n)\right] . \qquad (4)$$

Equation (4) can be computed efficiently using the real-time [37] finite-temperature [38] density matrix renormalization group [39, 40] (DMRG) algorithm introduced in [11]. DMRG is essentially controlled by the so-called discarded weight ϵ. We ensure that ϵ is chosen small enough and that N is chosen large enough to obtain numerically-exact results in the thermodynamic limit.

Non-equilibrium energy current — We start by studying a XXZ chain with two additional perturbations (dimerization $\lambda < 1$ and a staggered field $b > 0$) which both render the system nonintegrable [8, 30, 31]. At time $t = 0$, two semi-infinite chains each being prepared in thermal equilibrium at temperatures $T_{L,R}$ are coupled by h_0 to an overall translationally-invariant chain. Exemplary results for $\langle J_E(n,t)\rangle$ are shown in Figure 1. The current at the interface $n = 0$ saturates on a scale $t \sim 1$ [note the definition of units via Equation (2)] irrespective of the temperature difference $T_L - T_R$ or the absolute values of $T_{L,R}$ and regardless of the fact whether or



FIG. 3. (a) Temperature-dependence of the steady-state energy current (which becomes position-independent for $t \to \infty$) of the integrable XXZ chain in the gapless ($\Delta = 0.5$) and gapped ($\Delta = 2$) regime. The right temperature T_R is varied at fixed T_L. (b) The curves at different T_L collapse if shifted vertically; thus, the T-dependence is of the simple functional form $\lim_{t\to\infty}\langle J_E(n,t)\rangle = f(T_L)-f(T_R)$. The linear thermal conductance $\sim \partial_T f$ therefore completely determines the nonequilibrium current. A CFT approach [32] predicts universal behavior $f(T) = c\frac{\pi}{12}T^2$ at low T which we verify (Insets) for the gapless XXZ chain (central charge $c = 1$) and the quantum Ising model ($c = 1/2$). The free fermion case $\Delta = 0$ can be solved analytically [28, 29] at any T and provides a test for our DMRG numerics.

not the system is gapped. The only exception is $b > 0$ where $\langle J_E(n,t)\rangle$ does not reach a finite steady-state value within the time scales accessible by our numerics [Figure 1(c)], again irrespective of the fact whether or not b opens a gap. We will now try to understand this in more detail.

The time evolution of the local energy density $h(n,t) = \langle h_n(t)\rangle$ of the XXZ chain (which for a homogeneous system might be a measure for an effective temperature) is shown in the Inset to Figure 1(a). It does not reach a steady-state value but becomes increasingly smooth. This is not suprising since we are simulating a closed quantum system – but gives rise to the questions: (1) Why does the current saturate except for $b > 0$? (2) Are we actually modelling energy transport between reservoirs whose temperature is fixed [27]? Both are reasonable if it does not matter over which length scale L the temperature difference $T_L - T_R$ is applied; qualitatively, this should be the case if thermal transport properties of the chain are length-independent, i.e. if the thermal conductance G of a finite system does not decrease with its length L, or equivalently, if the conductivity $\sigma = GL$ of an infinite chain $L \to \infty$ is infinite. More quantitatively, we conjecture an intimate relation between nonequilibrium and linear response: The nonequilibrium energy current relaxes to a finite steady-state value if the linear-response thermal conductivity is infinite, i.e. if the Drude weight D is nonzero. To support this we now proceed by calculating D.

Linear response thermal Drude weight — The thermal Drude weight is determined by the long-time behavior of the energy current correlation function [2, 8],

$$D = \lim_{t\to\infty}\lim_{N\to\infty}\frac{\mathrm{Re}\,\langle J_E(t)J_E\rangle}{2NT^2}\ ,\ J_E = \sum_n J_E(n)\ , \quad (5)$$

which can be readily computed using DMRG. Results are shown in Figure 2. For $\lambda = 1$ and $b = 0$, J_E is conserved, thus $\langle J_E(t)J_E\rangle = \langle J_E(0)J_E\rangle$ [and the Drude weight can alternatively be obtained via Bethe ansatz [18]; see symbols in (a) and (b)]. The energy current correlation function of the dimerized chain seems to saturate at a finite value (or decays on a hidden large time scale). Our data thus indicates a nonzero Drude weight – which is interesting because the model is nonintegrable (previous results for D [8, 9, 15] are controversial). The deeper reason for this – the protection of the energy current by an unknown nonlocal conserved operator [3] – will be left as a subject for future work. Our data for $b > 0$, however, is consistent with $D = 0$ [see Figure 2(c)] both in the gapless and gapped regime. Recalling that the nonequilibrium current relaxes to a nonzero steady-state value in all cases where $b = 0$, the observation of a finite (vanishing) Drude weight for $\lambda \le 1$ ($b > 0$) supports our above conjecture.

Asymptotic current, homogeneous system — We now turn to study the temperature-dependence of the steady-state (position-independent) current. The result for the XXZ chain both in the gapless in gapped regime is illustrated in Figure 3(a). The asymptotic current seem to be of a strikingly simple functional form:

$$\lim_{t\to\infty}\langle J_E(n,t)\rangle = f(T_L) - f(T_R)\ , \quad (6)$$

indicating a second intimate relation between nonequilibrium and linear response: The linear thermal conductance $\sim \partial_T f(T)$ determines the steady-state nonequilibrium current at any $T_L - T_R$. Equation (6) can be established by varying T_R at fixed T_L; the corresponding curves collapse if shifted vertically [see Figure 3(b)]. The limiting behavior (both in the gapless and gapped regime) of $f(T)$ is given by

$$f(T) \sim \begin{cases} T^2 & T \ll 1 \\ T^{-1} & T \gg 1 \,. \end{cases} \tag{7}$$

Other details of f such as prefactors or the crossover scale (which for $\Delta = 0$ and $\Delta \gg 1$ is determined by the bandwidth or the size of the gap, respectively) in general depend on the model parameters. However, a recent conformal field theory approach [32] conjectures that the low-temperature behavior of a gapless system is universally given by $f(T) = c\frac{\pi}{12}T^2$ with c being the CFT central charge, which follows intuitively from the version of the Stefan-Boltzmann law satisfied by a CFT [33]. We confirm this prediction for the XXZ chain ($c = 1$) as well as the quantum Ising model ($c = 1/2$); this is illustrated in the Insets to Figure 3(b). This is a highly nontrivial result because: (1) It is unclear why for a microscopic model whose equilibrium physics is governed by a certain low-energy field theory the very same field theory should describe the long-time behavior of the microscopic model in nonequilibrium (note that the behavior for $0 < b < b_c$ is not captured by the CFT!), and (2) Even linear-response transport properties (such as the Drude weight) are not determined by the low-energy theory alone but by a delicate interplay of conserved quantities protecting the current and dangerously irrelevant operators scattering the latter [1–3].

Our results for the dimerized chain at $\lambda = 0.8$ are still consistent with Equation (6), indicating that it might be a universal property of any translationally-invariant system. At smaller λ and low T, we cannot reach time scales where oscillations of the current have died out completely; we thus cannot exclude a different behavior of models which are strongly nonintegrable.

The free fermion case $\Delta = 0$ can be solved exactly [28, 29]; Equation (6) reflects a noninteracting thermal Landauer-Büttiker formula. This analytic result can be used to test our DMRG numerics at any temperature [see the comparision in Figure 3(b) as well as in the Inset to Figure 4].

Inhomogeneous systems — We finally investigate systems which are not translationally invariant. We start by studying the effects of different bulk parameters $J_{n<0} = J_L$, $\Delta_{n<0} = \Delta_L$, $J_{n\geq 0} = J_R$, $\Delta_{n\geq 0} = \Delta_R$ (and $b_n = 0$). If these parameters are chosen such that the renormalized Fermi velocities in the left and right halves coincide, backscattering due to the barrier (which is naturally present at the interface $n = 0$ [42]) can be tuned to zero [41], and Luttinger liquid (LL) boundary physics [34] is absent. Results are displayed in Figure 4, indicating that Equation (6) might still hold.



FIG. 4. Steady state current for $\lambda = 1$, $b = 0$ but different bulk interactions $\Delta_L = 0.3, \Delta_R = 0.8$ and different couplings $J_L = 1, J_R = 0.808$ to tune the backscattering at the interface to zero [41]. The magnitude of the current is the same for $T_L = T_1, T_R = T_2$ and $T_L = T_2, T_R = T_1$. The curves at different T_2 collapes, indicating that Equation (6) might still hold. Inset: Linear thermal conductance of two identical XXZ chains connected by a strong barrier $J_0 = 0.002$. The low-T behavior is consistent with $T^{2/K-1}$ where K is the Luttinger liquid parameter [34]. At $\Delta = 0$ we repoduce the exact result of [29] for any T.

If two homogeneous XXZ chains $J_{n\neq 0} = 1, \Delta_n = \Delta$ are connected through a barrier J_0, the linear thermal conductance $G = \partial_{T_L}\langle J_E(t\to\infty)\rangle|_{T_L=T_R=T}$ is expected to feature a low-T power law $T^{2/K-1}$ with K being the LL parameter [34]. For $\Delta = 0$, Equation (6) still holds even if $J_0 < 1$ [29], and $2/K - 1 = 1$ just reflects the asymptotic T^2 behavior of f. Our data for G in presence of interactions is consistent with $T^{2/K-1}$. This is illustrated in the Inset to Figure 4 where J_0 is chosen small so that the scale on which LL boundary physics manifests becomes large [43]. Equation (6) still seems to hold above this scale.

Summary — In this paper we provided evidence for an intimate connection between nonequilibrium and linear-response thermal transport properties of isolated infinite spin chains (or equivalently, interacting spinless fermions): (1) The energy current of a system which initially features a temperature gradient $T_L \neq T_R$ saturates to a finite value if the equilibrium thermal Drude weight D is finite, and (2) The value of the steady-state current at arbitrary $T_{L,R}$ is completely determined by the linear thermal conductance. Our data suggests that $D > 0$ for a nonintegrable dimerized chain (or that the current correlation function decays on a hidden large time scale).

Acknowledgments — We are indebted to Ehud Altman, Fabian Heidrich-Meisner, Volker Meden, Kurt Schönhammer, and Dirk Schuricht for fruitful discussions and comments and acknowledge support by the Deutsche Forschungsgemeinschaft via KA3360-1/1 (C.K.) as well as by the AFOSR MURI on "Control of Thermal and Electrical Transport" (R.I.) and the Nanostructured Thermoelectrics program of LBNL (J.E.M. and C.K.).

---

[1] A. Rosch and N. Andrei, Phys. Rev. Lett. **85**, 1092 (2000).
[2] J. Sirker, R. G. Pereira, and I. Affleck, Phys. Rev. Lett. **103**, 216602 (2009).
[3] T. Prosen, Phys. Rev. Lett. **106**, 217206 (2011).
[4] B. S. Shastry and B. Sutherland, Phys. Rev. Lett. **65**, 243 (1990); X. Zotos, Phys. Rev. Lett. **82**, 1764 (1999); J. Benz, T. Fukui, A. Klümper, and C. Scheeren, J. Phys. Soc. Jpn. Suppl. **74**, 181 (2005).
[5] B. N. Narozhny, A. J. Millis, and N. Andrei, Phys. Rev. B **58**, R2921 (1998).
[6] J. V. Alvarez and C. Gros, Phys. Rev. Lett. **88**, 077203 (2002).
[7] D. Heidarian and S. Sorella, Phys. Rev. B **75**, 241104(R) (2007).
[8] F. Heidrich-Meisner, A. Honecker, D. C. Cabra, and W. Brenig, Phys. Rev. B **68**, 134436 (2003).
[9] F. Heidrich-Meisner, A. Honecker, and W. Brenig, Eur. Phys. J. Spec. Topics **151**, 135 (2007).
[10] P. Jung and A. Rosch, Phys. Rev. B **76**, 245108 (2007).
[11] C. Karrasch, J. H. Bardarson, and J. E. Moore, Phys. Rev. Lett. **108**, 227206 (2012).
[12] H. Castella, X. Zotos, and P. Prelovšek, Phys. Rev. Lett. **74**, 972 (1995).
[13] A. L. Chernyshyev, J. Magn. Magn. Matter. **310**, 1263 (2007).
[14] A. V. Sologubenko, K. Giannó, H. R. Ott, U. Ammerahl, and A. Revcolevschi, Phys. Rev. Lett. **84**, 2714 (2000).
[15] J. V. Alvarez and C. Gros, Phys. Rev. Lett. **89**, 156603 (2002).
[16] P. Jung, R. W. Helmes, and A. Rosch, Phys. Rev. Lett. **96**, 067202 (2006).
[17] X. Zotos, F. Naef, and P. Prelovšek, Phys. Rev. B **55**, 11029 (1997).
[18] A. Klümper and K. Sakai, J. Phys. A: Math. Gen **35**, 2173 (2002).
[19] K. Saito, Phys. Rev. B **67**, 064410 (2003).
[20] E. Orignac, R. Chitra, and R. Citro, Phys. Rev. B **67**, 134426 (2003).
[21] D. Gobert, C. Kollath, U. Schollwöck, and G. Schütz, Phys. Rev. E **71**, 036102 (2005).
[22] M. Michel, O. Hess, H. Wichterich, and J. Gemmer, Phys. Rev. B **77**, 104303 (2008).
[23] E. Boulat, H. Saleur, and P. Schmitteckert, Phys. Rev. Lett. **101**, 140601 (2008).
[24] S. Langer, F. Heidrich-Meisner, J. Gemmer, I. P. McCulloch, and U. Schollwöck, Phys. Rev. B **79**, 214409 (2009).
[25] T. Prosen and M. Žnidarič, J. Stat. Mech. 2009, **P02035** (2009).
[26] R. Steinigeweg, M. Ogiewa, and J. Gemmer, Europhys. Lett. 87, **10002** (2009).
[27] S. Ajisaka, F. Barra, C. Mejia-Monasterio, and T. Prosen, arXiv:1204.1321.
[28] L. Arrachea, G. S. Lozano, and A. A. Aligia, Phys. Rev. B **80**, 014425 (2009).
[29] L. Bruneau, V. Jaksic, C. Pillet , arXiv:1201.3190.
[30] This is supported by level statistics [31].
[31] Y. Huang, C. Karrasch, and J. E. Moore, unpublished.
[32] D. Bernard and B. Doyon, J. Phys. A **45**, 362001 (2012).
[33] J. Cardy, J. Stat. Mech. 2010, **P10004**, 2010.
[34] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **76**, 3192 (1996).
[35] F. D. M. Haldane, Phys. Lett. **81A**, 153 (1980).
[36] F. C. Alcaraz and A. L. Malvezzi, J. Phys. A: Math. Gen. **28**, 1521 (1998).
[37] G. Vidal, Phys. Rev. Lett. **93**, 040502 (2004); S. R. White and A. E. Feiguin, Phys. Rev. Lett. **93**, 076401 (2004); A. Daley, C. Kollath, U. Schollwöck, and G. Vidal, J. Stat. Mech.: Theory Exp. P04005 (2004); P. Schmitteckert, Phys. Rev. B **70**, 121302(R) (2004).
[38] A. E . Feiguin and S. White, Phys. Rev. B **72**, 220401(R) (2005); T. Barthel, U. Schollwöck, and S. R. White, Phys. Rev. B **79**, 245101 (2009).
[39] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).
[40] U. Schollwöck, Ann. Phys. **326**, 96 (2011).
[41] N. Sedlmayr, J. Ohst, I. Affleck, J. Sirker, and S. Eggert, Phys. Rev. B **86**, 121302(R) (2012).
[42] K. Janzen, V. Meden, and K. Schönhammer, Phys. Rev. B **74**, 085301 (2006).
[43] V. Meden, S. Andergassen, W. Metzner, U. Schollwöck, and K. Schönhammer, Europhys. Lett. 64, **769** (2003).